

Janet M. Intrieri¹ and Matthew D. Shupe²¹NOAA/Environmental Technology Laboratory, Boulder, Colorado²Cooperative Institute for Research in the Environmental Sciences, NOAA/University of Colorado, Boulder, Colorado

1. INTRODUCTION

Clouds play an especially important and amplified role at high latitudes because the polar surface is characterized by high albedos and the polar atmosphere is much drier than lower latitudes. Arctic clouds also strongly influence the yearly sea-ice cycle of melting and formation so any potential feedback from clouds, in a changing climate scenario, can be substantially important. Changes in surface temperature, sea-ice concentration, atmospheric stability, etc. will undoubtedly affect the properties of clouds, and a change in cloud properties will, in turn, affect the radiation at the sea-ice surface. However, neither this cloud-radiation feedback mechanism, nor is its relation to the ice-albedo feedback, both critical components in a changing climate scenario at high latitudes, are well understood.

In this study, we utilize Arctic datasets from two distinctly different Arctic regions to a) create a baseline of data in a region where measurements are generally lacking, b) investigate the differences between coastal Alaskan and Arctic Ocean region clouds, and c) determine the respective influences of clouds on the surface radiation budget. The cloud and radiation datasets were obtained at the DOE's Atmospheric Radiation Measurement (ARM) North Slope of Alaska (NSA) facility in Barrow, Alaska, from 1998 to 2003, and the Surface Heat Budget of the Arctic (SHEBA) program, obtained over the Western Arctic Ocean region during 1997-1998. Similar cloud and radiation observations obtained in both locations by lidar, radar, and radiometers allow for a comparison of cloud occurrence and physical properties between these two distinctly different regions and were used to produce annual cycles of cloud occurrence and height, atmospheric temperature and humidity, surface broadband fluxes, surface albedo, and cloud radiative forcing.

2. INSTRUMENT AND ANALYSIS INFORMATION

The instrumentation from the SHEBA field program and the NSA facility used as part of this study were similar with only one notable difference: the lidar at SHEBA, the Environmental Technology Laboratory's (ETL) Depolarization and Backscatter Unattended Lidar (DABUL), included depolarization ratio measurements. Depolarization ratio data provide a means for assessing

the presence of liquid versus ice phase in a cloud layer. (We note that lidar depolarization measurements have recently been added at the NSA). Otherwise, the measurements were ostensibly identical and included cloud reflectivity from the Millimeter Cloud Radar (MMCR), shortwave (SW) and longwave (LW) fluxes from upward and downward looking broadband hemispheric radiometers, liquid water path (LWP) derived from microwave radiometer, and temperature and humidity profiles from atmospheric soundings. Although, the soundings at SHEBA and NSA provide identical information, the frequency of the soundings was much greater during SHEBA (2-4 per day/everyday) versus those obtained at NSA (once per day on weekdays only). Cloud morphology information such as frequency of occurrence are created from a combination of lidar, ceilometer, and radar data to provide the best possible estimate. For additional information on SHEBA instruments please refer to Intrieri et al. (2002a) and Persson et al. (2002). For additional information on the NSA site and instrument specifics refer to www.arm.gov.

Annual cycles of surface shortwave and longwave fluxes and surface cloud radiative forcing (SCF), defined as the difference between the all-sky and clear-sky net surface radiative fluxes, have already been produced for SHEBA (Intrieri et al. 2002b). Cloud forcing provides an estimate of how much a cloud changes the surface radiative fluxes relative to clear skies, and has become a common means for quantifying the radiative relationships between clouds and the radiation budget (Ramanathan et al. 1989). Definitions for the LW, SW and total SCF's, are given as:

$$CFLW = F(Ac) - F(0) \quad (1)$$

$$CFSW = Q(Ac) - Q(0) \quad (2)$$

$$SCF = CFLW + CFSW \quad (3)$$

Where, Ac is the cloud fraction and F and Q are the net surface LW and SW fluxes, respectively. Positive forcing values indicate that clouds impart a warming effect at the surface relative to clear skies (i.e., a greenhouse effect) and negative forcing values indicate that clouds impart a cooling effect relative to clear skies (i.e., the albedo effect). Cloud forcing at the surface is calculated using ground-based measurements of broadband fluxes for the all-sky values and modeled fluxes for the clear-sky values since clear skies occur so infrequently in the Arctic. In order to derive the SHEBA and NSA surface cloud forcing values, modeled clear-sky radiation results were calculated using the Santa Barbara DISORT Atmospheric Radiative Transfer code (SBDART: Richiazzi et al. 1998). The NSA fluxes and SCF's were

Corresponding author address: Janet Intrieri, NOAA/ Environmental Technology Laboratory, 325 Broadway, Boulder, CO 80305; e-mail: janet.intrieri@noaa.gov

calculated and processed in the same way as the SHEBA dataset for consistency and to minimize error.

Measurements used for input into the SBDART model included temperature and relative humidity profiles from radiosondes and surface albedo from the SW radiometer. Interpolating atmospheric conditions between sounding times can introduce errors given that weather can change significantly and weather systems can pass through undetected in that amount of time. During SHEBA, radiosondes were launched every 6-12 hours (7 days per week) therefore, interpolating between times introduced less error than interpolating between times at NSA where soundings were only obtained once per day on weekdays. The surface albedo, a critical parameter in cloud forcing calculations, was calculated from the upward and downward looking SW radiometer hourly averaged flux values at both SHEBA and NSA. These measurements are therefore most representative of the local areas within view of the radiometers.

3. SHEBA AND NSA SURFACE CLOUD FORCING

3.1 Total Cloud Forcing Annual Cycle

The annual cycle of total SCF (CFLW + CFSW) at SHEBA (Fig. 1, top panel – solid bold line) shows that clouds induced surface warming throughout most of the year with only a short period of surface cooling in the middle of summer, when cloud shading effects overwhelmed cloud greenhouse effects. At NSA, the annual trend is the same, (warming in winter, spring and fall), but the summer cooling effect is much stronger and lasts much longer when compared to SHEBA. The total SCF at NSA falls below 0 Wm^{-2} , dipping down to $\sim -100 \text{ Wm}^{-2}$, for 3 months (from early June to mid September) while at SHEBA, the total SCF only fell below 0 Wm^{-2} , dipping to -5 Wm^{-2} , for the month of July. The seasonal averages for the total SCF at SHEBA are 30 Wm^{-2} for fall, winter and spring and -10 Wm^{-2} for summer (defined as the time period when the SCF is negative). At NSA, the seasonal averages (averaged over all six years) are 37 Wm^{-2} for fall, winter and spring and -60 Wm^{-2} during summer (defined same as above).

The greatest year-to-year variability at NSA is noted during the summer melt season and the least amount of year-to-year variability is observed during the fall freeze-up. The yearly variability of the total SCF at NSA between the six years of data is $\sim 30 \text{ Wm}^{-2}$ during winter, spring and fall, however, during early summer, the variability is greater, varying by as much as $\sim 80 \text{ Wm}^{-2}$ between 1998 and 2002. The variability in magnitude and the timing of the total SCF is greatest during the early summer season due to fluctuations in the timing of the melt season onset and local weather-related events that produce extended periods of clear skies and fluctuations in cloud amount and precipitation (which affects albedo). By the same reasoning, the data suggest that the fall season shows a much more consistent cloud and weather pattern (as well as albedo remaining high with lowering temperatures) between each NSA year, translating to greater consistency in

cloud forcing. Fig. 2 illustrates 20-day averages of cloud fraction from NSA and SHEBA. Note the more consistent values between day 250 and 320 (September through November) with cloud fractions varying only between 85 and 100% whereas between day 150 and 250 (June through August) the fractions fluctuate between 70 and 95%.

In general, the annual cycle of NSA total cloud forcing reveals several points: 1) cloud amounts are fairly consistent and did not change substantially in the Barrow region during the fall, winter and spring seasons over the six year analysis period, 2) late spring and summer are most variable since there are more late spring season weather-related disturbances that vary the onset of the melt season, and 3) cloud's in the Barrow region have fairly consistent boundaries by which they affect the surface radiation ($< 50 \text{ Wm}^{-2}$ in fall, winter and spring and $> -150 \text{ Wm}^{-2}$ in summer). In order to understand the specific differences between the two regional annual cycles, including the inter-annual melt season onset variability and smaller-scale anomalous events at NSA, we discuss the individual LW and SW component parts below.




Fig. 1. Annual cycle of 20-day average a) Total, b) LW, and c) SW surface cloud forcing for SHEBA (bold line) plus six years of NSA data. All values in Wm^{-2} .

3.1 Longwave Cloud Forcing Annual Cycle

Cloud and radiation study results from SHEBA (Shupe and Intrieri 2004) showed that LW Cloud Forcing (CFLW) is a function of cloud temperature, height, and emissivity (which is a function of cloud microphysical properties). The trend and magnitude of the cloud warming effect on the surface during the entire annual cycle is fairly similar at both SHEBA and NSA, with clouds warming the surface in the LW between 10 and 70 Wm^{-2} throughout the year (Fig. 1, middle panel). This warming is predominantly due to the fact that clouds are optically thicker (i.e. stronger emitters) than the clear atmosphere and the warming is accentuated by the frequent occurrence of temperature inversions that often cause clouds to emit at temperatures warmer than the surface. In winter, surface temperatures are much warmer under cloudy skies than during clear

conditions and in all seasons clouds containing small liquid water droplets are efficient emitters in the longwave, becoming essentially black at LWP's greater than 30 gm^{-2} . The average LWCF during SHEBA was $\sim 38 \text{ Wm}^{-2}$ while at NSA the 6-year averages were $\sim 45 \text{ Wm}^{-2}$.

The similarity in the magnitudes of the CFLW between the two regions is due to the similar atmospheric temperature structure and cloudiness trends found at both locations. The differences in the LWCF are attributed to the relative amount of clouds observed between the two sites. In general, NSA had greater cloud occurrence values in all seasons except for summer than was observed at SHEBA (Fig. 2). For example, in early winter at SHEBA (bold line between days 0 and 50) there were substantially less clouds present than at NSA so the forcing values are smaller. During mid-summer (\sim Day 200), for all years except 2003, NSA was less cloudy than SHEBA, so the LW forcing values are accordingly lower during that period at NSA. By the same logic, there is a fall season (September-November: \sim days 240 and 310) CFLW maximum present in all of the measurement years which coincides with the cloudiness occurrence maximum values in both locations. In general, the CFLW response is positive and fairly flat throughout the year so, in and of itself, the longwave component does not introduce the significant amount of seasonal structure present in the total SCF.




Fig. 2. 20-day averages of cloud fraction for SHEBA (solid line) and six years of data from NSA.

3.3 Shortwave Cloud Forcing Annual Cycle

Cloud and radiation results from SHEBA (Shupe and Intrieri 2004) showed that SW Cloud Forcing (CFSW) is a function of cloud transmittance (i.e., cloud microphysical properties), surface albedo, and the solar zenith angle. The CFSW cloud cooling effect (i.e., the SW shading effect of clouds) in summer is significantly stronger at NSA than at SHEBA (Fig. 1, bottom panel). At SHEBA, the 20-day average CFSW values dipped to $\sim -50 \text{ W m}^{-2}$ in July (centered around day 190). At NSA, the 20-day average CFSW values typically ranged two

to three times greater than the SHEBA values, dipping to more negative values between -100 to -150 W m^{-2} .

This amplified shortwave radiative effect during summer at NSA is directly related to the much lower surface albedos (Fig. 3) as well as the somewhat higher insolation values and Sun angles. Generally speaking, the snow/ice surface never fully melted away at SHEBA. For practical purposes, the radiometers at SHEBA were sited on a multi-year ice floe so that the instrumentation wouldn't need to be relocated due to melting. In fact, ablation shields (plywood boards painted white) were placed around the base of the meteorological tower to ensure that the instruments would make it through the summer. In this manner, the albedos calculated at the singular radiometer point near the tower were kept artificially higher with typical albedos being 0.10-0.15 greater than the average albedos determined from a 200 m line (see Fig. 5, Intrieri et al. 2002b).




Fig. 3. Annual cycles of albedo for SHEBA (solid line) and six years of data from NSA.

Taking into account the lower line-averaged values at SHEBA produced lower total cloud forcing values, dipping below 0 W m^{-2} between early June to mid-August and a minimum value of $\sim -45 \text{ W m}^{-2}$ (see Fig. 11, Intrieri et al. 2002b). However, even at these lower SHEBA albedo values, the NSA SW and total cloud forcing are still twice as large. The reason boils down to the fact that at NSA the snow fully melts away to reveal bare dirt and tundra. At SHEBA, the minimum albedo values were 0.50 for the point measurement and 0.38 for the line-averaged measurement, while at NSA these values are typically ~ 0.15 . Together, those data indicate that the surface albedo is the major source of CFSW difference between SHEBA and NSA.

Solar Zenith Angle (SZA) also contributes to the differences observed between the CFSW at SHEBA and NSA. The SZA determines the potential amount of solar radiation available at the surface and how long the sun is above the horizon. Since the SHEBA drift site was $\sim 1000 \text{ km}$ North of Barrow, the minimum summer SZA was 54° compared to a minimum SZA of 50° at Barrow. Thus, at SHEBA the annual maximum insolation was about 650 W m^{-2} , while the NSA maximum is nearly 750 W m^{-2} . Shupe and Intrieri (2004) found that at SHEBA the SWCF was always less than 25%, but typically around 3-10%, of the total insolation. Therefore, a small but significant portion of the CFSW difference between SHEBA and NSA can be directly attributed to SZA.

As stated above, some of the year to year variability at the NSA site is directly related to the onset of the melt season and its impact on surface albedo. On average, the melt onset occurs around June 9 to 19 (day 160-170) but was observed as early as May 27 (day 147) in 2002 and as late as June 28




Fig. 4 . Scatterplot of CFLW (top) and CFSW (bottom) versus cloud fraction amount for NSA, 2002.

(day 179) in 1999. After melt has fully occurred the variability in the observed CFSW is often due to extended periods of clear skies present over Barrow Alaska (Fig. 1, bottom panel; e.g. ~day 190 during 1998, 2001, 2002) when the forcing values effectively increase because forcing under clear skies is zero. In spite of the variable spring melt onset, there appears to be little variability in the timing of the fall transition back to a snow covered surface indicating that once snow falls, it remains cold enough to keep the surface albedo high.

3.4 Sensitivity of Cloud Forcing

We examined the sensitivity of the cloud occurrence to the cloud forcing for both the LW and the SW forcing values. By definition, cloud forcing under clear skies is equal to zero and under 100% overcast skies the forcing is at its maximum. At SHEBA, we found that the sensitivity of CFSW to cloud fraction was negatively correlated and ranged between 0.0 and -1.0 W m^{-2} per percent cloudiness (Shupe and Intrieri 2004) whereas at NSA values approached twice that to around -2.0 W m^{-2} per percent cloudiness under overcast sky conditions (Fig. 4). At SHEBA, the CFLW and cloud fraction were positively correlated and the sensitivity was about 0.65 W m^{-2} per percent cloudiness. At NSA the sensitivity is slightly higher at about 0.75 W m^{-2} per percent cloudiness (Fig. 4). In summary, the warming effect of clouds on the surface during winter is similar at SHEBA and NSA, mostly due to the ever-present inversion and similar atmospheric temperatures, while the summer shading effect of clouds at Barrow is much greater than at SHEBA because of the significantly lower surface albedos and higher insolation values.

4. SUMMARY AND FUTURE WORK

Cloud and radiation data from two distinctly different Arctic areas are analyzed to study the differences between coastal Alaskan and open Arctic Ocean region clouds and their respective influence on the surface radiation budget. Radar, lidar, radiometer, and sounding measurements from both locations were used to produce annual cycles of cloud properties, atmospheric temperature and humidity, surface longwave and shortwave broadband fluxes, surface albedo, and cloud radiative forcing.

In general, both regions revealed a similar annual trend of cloud occurrence fraction with minimum values in winter (60-75%) and maximum values during spring, summer and fall (80-90%). However, the annual average cloud occurrence fraction for SHEBA was lower (76%) than the 6-year average cloud occurrence at NSA (92%). Both Arctic areas also showed similar annual cycle trends of cloud forcing with clouds warming the surface through most of the year and a period of surface cooling during the summer, when cloud shading effects overwhelm cloud greenhouse effects. The greatest difference between the two regions was observed in the magnitude of the cloud cooling effect (i.e., shortwave cloud forcing), which was significantly stronger at NSA and lasted for a longer period of time than at SHEBA. This is predominantly due to the longer and stronger melt season at NSA (i.e., albedo values that are much lower coupled with Sun angles that are somewhat higher) than the melt season observed over the ice pack at SHEBA. Longwave cloud forcing values were comparable between the two sites indicating a general similarity in cloud cycle and atmospheric temperature and humidity structure between the two regions.

Acknowledgments. This research was supported by the Biological and Environmental Research Program (BER), U.S. Department of Energy, Interagency Agreement DE-AI03-02ER63325, the NSF SHEBA Agreement OPP-9701730, and the NASA FIRE-ACE Contract L64205D.

5. REFERENCES

- Intrieri, J.M., and M. D. Shupe, 2004: Characteristics and radiative effects of diamond dust over the Western Arctic Ocean region. *J. Climate*, **17**, 2953-2960.
- , M. D. Shupe, B. J. McCarty, T. Uttal, 2002a: Arctic cloud statistics from lidar and radar at SHEBA. *J. Geophys. Res.*, **107**, 8030.
- , C. F. Fairall, O. G. P. Persson, M. D. Shupe, and R. M. Moritz, 2002b: Annual cycle of cloud forcing over the Arctic. *J. Geophys. Res.*, **107**, 8039.
- Ramanathan V., et al., 1989: Climate and the Earth's radiation budget. *Physics Today*, 22-32.
- Ricchiazzi, P., S. Yang, C. Gautier, and D. Sowle, 1998: SBDART: A research and teaching software tool for plane-parallel radiative transfer in the Earth's atmosphere. *Bull. Am. Meteor. Soc.*, **79**, 2101-2114.
- Shupe, M. D., and J. M. Intrieri, 2004: Cloud radiative forcing of the Arctic Surface: The influence of cloud properties, surface albedo, and solar zenith angle. *J. Climate*, **17**, 616-628.